

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd

Document heading

doi: 10.1016/S2222-1808(14)60733-7

© 2014 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

Molecular study on diarrheagenic *Escherichia coli* pathotypes isolated from under 5 years old children in southeast of IranHesam Alizade¹, Reza Ghanbarpour², Mohammad Reza Aflatoonian^{3*}¹Research Center for Tropical and Infectious Diseases, Kerman University of Medical Sciences, Department of Microbiology, Sirjan Faculty of Medical Sciences, Kerman University of Medical Sciences, Kerman, Iran²Molecular Microbiology Department, Faculty of Veterinary Medicine, Shahid Bahonar University, Zoonosis Research Committee of Kerman University of Medical Sciences, Kerman, Iran³Leishmaniose Research Committee of Kerman University of Medical Sciences, Research Center for Tropical and Infectious Diseases, Kerman University of Medical Sciences, Kerman, Iran

ARTICLE INFO

Article history:

Received 12 May 2014

Received in revised form 27 May 2014

Accepted 22 Aug 2014

Available online 28 Aug 2014

Keywords:

Escherichia coli

Diarrhea

Phylogenetic group

ABSTRACT

Objective: To determine the phylogenetic groups and prevalence of diarrheagenic *Escherichia coli* (*E. coli*) (DEC) genes from children less than five years of age with diarrhea in southeast of Iran.**Methods:** A total of 142 *E. coli* isolates were isolated from diarrheic samples. The isolates were examined for detection of virulence determinants and their phylogenetic background by PCR technique.**Results:** The *E. coli* isolates fall into four phylogenetic groups: A (40.14%), B1 (18.31%), B2 (16.90%) and D (24.65%). Eighty isolates were positive for at least one of the examined DEC genes. *E. coli* isolates were classified in enterotoxigenic *E. coli* (52 isolates), enteroaggregative *E. coli* (23), atypical enteropathogenic *E. coli* (9), enteroinvasive *E. coli* (2).**Conclusions:** This study demonstrated the importance of enterotoxigenic *E. coli* and enteroaggregative *E. coli* pathotypes in the childhood diarrhea. An epidemiologic surveillance especially for DEC, would be useful in control and prevention of infectious diarrhea in children.

1. Introduction

Gastrointestinal infections due to pathogenic *Escherichia coli* (*E. coli*) are significant causes of morbidity and mortality in children, particularly in developing countries[1]. Clinical categories of *E. coli* comprise commensal, intestinal pathogenic and extra-intestinal pathogenic strains. Diarrheagenic *E. coli* (DEC) pathotypes include enterotoxigenic *E. coli* (ETEC), enteroaggregative *E. coli* (EAggEC), enteroinvasive *E. coli* (EIEC), enteropathogenic *E. coli* (EPEC), enterohemorrhagic *E. coli* (EHEC) and diffusely

adherent *E. coli*[2]. ETEC pathotype defined by the presence of plasmid-encoded enterotoxins, comprise thermostable toxin (*ST*) and the thermolabile toxin (*LT*). ETEC strains are the most common cause of childhood diarrhea among all *E. coli* pathotypes and the major cause of diarrhea in travelers to developing countries[3]. Several virulence factors of EAggEC associated with diarrhea in children. Most of the genes encoding these virulence factors are located in the pAA plasmid, such as probe CVD432 and transcriptional factor encoded by the *aggR* gene. The pAA plasmid also carries the *aap* gene, which secreted low-molecular weight protein that promotes dispersal of EAggEC on the intestinal mucosa and facilitates efficient colonization[4,5]. Outbreaks of EIEC diarrhea are usually food or water-borne. However, through person-to-person transmissions have also been reported[6]. EIEC strains are able to attack intestinal epithelial cells. The invasion plasmid antigen H

*Corresponding author: Mohammad Reza Aflatoonian, Leishmaniose Research Committee of Kerman University of Medical Sciences, Research Center for Tropical and Infectious Diseases, Kerman University of Medical Sciences, Kerman, Iran.

Tel: +98-913398 2084

Fax: +98-341211 2794

E-mail: alizade.h2000@yahoo.com

Foundation Project: Supported by a grant (No.: 111.IDT.91) from Research Center for Tropical and Infectious Diseases, Kerman University of Medical Sciences, Kerman, Iran.

(*ipaH*) gene sequence is used for the diagnosis of EIEC[7,8]. EPEC strains express *eaeA* gene, which produce intimin, and bundle forming pili (*bfpA*) responsible for the attaching and effacing lesions of intestinal microvilli[3,9]. Shiga-toxin-producing *E. coli* or EHEC are principal emerging pathogens that cause food and water-borne diarrheal diseases in humans. All Shiga-toxin-producing *E. coli* strains possess *stx1* and/or *stx2* genes that produce two powerful cytotoxins, called Shiga toxin[10]. The *eaeA* gene of EHEC shares considerable homology with the *eaeA* gene of EPEC. Attaching and effacing *E. coli* strains (*eaeA*+) that harbor the *bfpA* gene are classified as typical EPEC and strains that do not possess *bfpA* gene are classified as atypical EPEC[11,12]. There are important regional differences in the prevalence of different categories of DEC in South and Southeast Asia[13].

Strains of the phylogenetic groups differ in their genotypic and phenotypic characteristics, comprising their antibiotic-resistance profiles, their ability to exploit different sugars sources and their growth rate temperature relationships. Phylogenetically, *E. coli* strains are divided upon amplification of *chuA* and *yjaA* genes and DNA fragment TSPE4.C2. The patterns of amplicons assigned four groups A, B1, B2 and D. DEC strains are derived from groups A, B1 and D, non-pathogenic commensal strains from A and B1, and extra-intestinal pathogenic strains usually belong to groups B2 and D[2,14].

The purpose of this study was to analyze the distribution of phylogenetic groups and occurrence of diarrheagenic genes in *E. coli* isolated from children less than five years of age with diarrhea in southeast of Iran by PCR.

2. Materials and methods

2.1. Sampling and bacteriological identification

One hundred and forty two *E. coli* isolates were obtained from diarrheal samples of children under five years old. Isolates were collected between 2010 and 2012 from children referring to the laboratories of Kerman Province, southeastern Iran. Samples were cultured on Mac Conkey agar and eosin methylene blue (Biolife Laboratories, Milano, Italy). Standard bacteriological methods were used to confirm the *E. coli* isolates. Isolates were stored in Luria-Bertani broth (Invitrogen, Paisley, Scotland) with 30% sterile glycerol at -70 °C for further analysis.

2.2. Reference strains

Five *E. coli* strains were used as positive controls: *E. coli* H10407 for ETEC (*LT*+, *ST*+), *E. coli* 85b for EIEC (*ipaH*+

E. coli O42 for EAggEC (probe CVD432+, *aggR*+ and *aap*+), *E. coli* Sakai for EHEC and atypical EPEC (*stx1*+, *stx2*+ and *eaeA*+) and *E. coli* ECOR62 for (*chuA*+, *yjaA*+ and TspE4.C2+). *E. coli* strain MG1655 was used as a negative control for virulence genes. All the reference strains were from the bacterial collection of Microbiology Department of Ecole Nationale Vétérinaire Toulouse, France.

2.3. PCR protocol

DNA was extracted from *E. coli* isolates and reference strains by lysis method. All isolates were tested by multiplex PCR assay for the presence of the *LT*, *ST* and *ipaH* genes by Aranda *et al.*[4], for *stx1*, *stx2* and *eaeA* genes by China *et al.* and probe CVD432, *agg*, *aap* genes by Cerna *et al.*[15,16]. The phylogenetic groups (A, B1, B2, and D) of each *E. coli* isolate were carried out by triplex PCR method as described previously[17]. The primers used for detecting sequences encoding virulence genes and phylogenetic groups are described in Table 1.

Table 1
Oligonucleotide primers used in this study.

	Gene or probe name	Primer sequence (5'–3')	Product size (bp)	Reference
ETEC	<i>LT</i>	GGC GAC AGA TTA TAC CGT GC CGG TCT CTA TAT TCC CTC TT	450	[4]
	<i>ST</i>	ATT TTT CTT TCT GTA TTG TCT T CAC CCG GTA CAA GCA GGA TT	190	
EAggEC	Probe CVD432	CTG GCG AAA GAC TGT ATC AT CAA TGT ATA GAA ATC CGC TGT T	600	[16]
	<i>aggR</i>	CTA ATT GTA CAA TCG ATG TA AGA GTC CAT CTC TTT GAT AAG	457	
	<i>aap</i>	CTT GGG TAT CAG CCT GAA TG AAC CCA TTC GGT TAG AGC AC	310	
EIEC	<i>ipaH</i>	GTT CCT TGA CCG CCT TTC CGA TAC CGT C GCC GGT CAG CCA CCC TCT GAG AGT AC	600	[4]
EPEC & EHEC	<i>eaeA</i>	AGG CTT CGT CAC AGT TG CCA TCG TCA CCA GAG GA	570	[15]
	<i>stx1</i>	AGA GCG ATG TTA CGG TTT G TTG CCC CCA GAG TGG ATG	388	
	<i>stx2</i>	TGG GTT TTT CTT CGG TAT C GAC ATT CTG GTT GAC TCT CTT	807	[17]
Phylo-group	<i>yjaA</i>	TGA AGT GTC AGG AGA CGC TG ATG GAG AAT GCG TTC CTC AAC	211	
	TspE4.C2	CTG GCG AAA GAC TGT ATC AT CGC GCC AAC AAA GTA TTA CG	152	
	<i>chuA</i>	GAC GAA CCA ACG GTC AGG AT TGC CGC CAG TAC CAA AGA CA	279	

3. Results

3.1. Phylogenetic grouping

The triplex PCR assays for phylotyping of isolates revealed that isolates fall into four phylogenetic groups, whereas 40.14% (57 isolates) belonged to A, 18.31% (26 isolates) to B1, 16.90% (24 isolates) to B2 and 24.65% (35 isolates) to D phylogenetic groups.

3.2. Detection of DEC isolates

Multiplex PCR were performed to detect the main five categories of *E. coli*. PCR assays revealed that 80 isolates were positive for at least one of the examined DEC genes. Fifty two (36.62%) isolates were positive for *LT* and/or *ST* genes. The ETEC pathotype coding genetic marker *ST* and *LT* were the most prevalent genes in the isolates, while were detected in 11.97% and 9.86% of isolates respectively. Among 52 isolates possess ETEC pathotype genes 21 isolates (14.79%) were positive for both *LT* and *ST* genes (Table 2). Overall 23 (16.20%) of the 142 *E. coli* isolates analyzed carried the EAggEC encoding genes, while probe CVD432 and *aap* genes were detected in 9.86% and 6.34% of isolates respectively. None of the isolates were positive for *aggR* gene (Table 2). Of the 142 isolates investigated, nine (6.34%) isolates were positive for atypical EPEC pathotype coding genetic marker *eae*. Out of *E. coli* isolates analyzed 2 (1.41%) isolates had the gene genetic marker for *ipaH*, which characterized as EIEC pathotype. Of the all isolates surveyed, none were positive for the EHEC encoding genes (*stx1* and *stx2*) (Table 2).

Table 2

Distribution of pathotypes in phylogenetic groups from children less than five years old.

DEC	Gene	Total No. (%)	Phylo-group			
			A	B1	B2	D
ETEC	<i>LT</i>	14 (9.86)	—	7 (50.00)	5 (35.71)	2 (14.29)
	<i>ST</i>	17 (11.97)	5 (29.41)	7 (41.18)	—	5 (29.41)
	<i>LT/ST</i>	21 (14.79)	17 (80.96)	2 (9.52)	2 (9.52)	—
EAggEC	Probe CVD432	14 (9.86)	—	—	5 (35.71)	9 (64.29)
	<i>aap</i>	9 (6.34)	2 (22.22)	—	2 (22.22)	5 (55.56)
	<i>aggR</i>	—	—	—	—	—
EIEC	<i>ipaH</i>	2 (4.92)	—	—	—	2 (100.00)
EPEC	<i>eaeA</i>	9 (6.34)	2 (22.22)	—	7 (77.78)	—
EHEC	<i>stx1</i>	—	—	—	—	—
	<i>stx2</i>	—	—	—	—	—
Total		86 (60.56)	26 (18.30)	16 (11.26)	21 (14.79)	23 (16.20)

3.3. Distribution of DEC genes in phylo-groups

Among 142 *E. coli* isolates 56.34% (80 isolates) and 43.66% (62 isolates) were positive and negative for at least one of the examined DEC genes respectively which, distributed in four phylo-groups (Table 3). ETEC strains were present among the isolates from A (21 isolates), B1 (17 isolates), B2 (7 isolates) and D (7 isolates) phylogenetic groups. Fourteen *LT* positive isolates belonged to B1 (7 isolates), B2 (5 isolates) and D (2 isolates) phylogenetic groups, while 17 isolates possess *ST* gene segregated in phylogenetic group A (5 isolates), B1 (7 isolates) and D (5 isolates). Phylotyping of *LT/ST* positive isolates showed that the isolates belonged to A (17 isolates), B1 (2 isolates) and B2 (2 isolates) phylo-groups. EAggEC strains encoding probe CVD432 fell into B2 (5 isolates) and D (9 isolates) phylogenetic groups. The *aap* positive isolates

were distributed in A (2 isolates), B2 (2 isolates) and D (5 isolates) phylogenetic groups. The atypical EPEC isolates were segregated in A (2 isolates) and B2 (7 isolates) phylo-groups. The EIEC strains coding genetic marker *ipaH* belonged to D (2 isolates) phylogenetic group (Table 2).

Table 3

The positive and negative isolates for at least one of the examined DEC genes distribute in phylo-groups.

	Phylo-group No. (%)				Total
	A	B1	B2	D	
Positive	24 (30.00)	16 (20.00)	16 (20.00)	24 (30.00)	80 (100.00)
Negative	33 (53.23)	10 (16.13)	7 (11.29)	12 (19.35)	62 (100.00)
Total	57 (40.14)	26 (18.31)	23 (16.20)	36 (25.35)	142 (100.00)

4. Discussion

DEC is recognized as an important cause of both outbreaks and sporadic cases throughout the world. There are at least six pathotypes of *E. coli* including ETEC, EAggEC, EIEC, EPEC, EHEC and diffusely adherent *E. coli*, which can cause intestinal infection in humans. Phylogenetic analysis of *E. coli* isolates showed that DEC strains were distributed among groups A, B1 and D and commensal strains in groups A and B1[14]. On the other hand, surveying the evolutionary origins of pathogenic *E. coli* is to determine the phylogeny distribution of the virulence genes[18]. DEC are second most common cause of diarrhea among children under five years old[3].

The results of the present study highlight the importance of ETEC as a cause of childhood diarrhea in the studied region of Kerman, Iran. ETEC is the major etiologic agents but under-recognized bacterial cause of either infantile diarrhea in all age groups in areas with poor sanitation. This pathotype is the most important cause of traveler's diarrhea; the organism is regularly imported to the developed world[19,20]. According to the results *ST*+ and *LT*+ isolates were detected in 11.97% and 9.86% of isolates respectively. In the other parts of world, there were reports differences from prevalence of ETEC pathotype. In studies on capital of Iran (Tehran) and Nicaragua 6.73% and 20.5% of diarrheic isolates obtained from children were positive for ETEC pathotype respectively[21,22]. Perez *et al.* indicated that 7.69% of *E. coli* isolates possessed ETEC encoding sequences[3]. In this study PCR results of phylogenetic determination, showed that ETEC pathotype mostly fell into group A, followed by B1, B2 and D. Escobar-Paramo *et al.* indicated that ETEC strains were found in A and B1 phylogenetic groups[23]. In a study, distribution of ETEC strains in phylo-groups were B1, A and D[3]. In the current study EAggEC pathotype encoding genes were examined. According to the results, probe CVD432+ and *aap*+ isolates were detected in 9.86% and 6.34%

of isolates respectively. The EAggEC pathotype has been implicated in endemic diarrhea among children in both industrialized and resource-poor countries[24]. In Tanzania a study on EAggEC isolates obtained from children less than five years old showed that prevalence of *aggR*+, *aap*+ and *astA*+ isolates were 61.6%, 26.7% and 15.1% respectively[5]. In Romanian, 11.6% of diarrheic isolates were positive for EAggEC pathotype that segregated to A (19 isolates), B1 (2 isolates), B2 (5 isolates) and D (3 isolates) phylogenetic groups[25], whereas in the current study 16.20% of isolates were positive for EAggEC pathotype and belonged to A, B2 and D phylo-groups. Boisen *et al.* surveyed potential virulence factors among 121 EAggEC strains isolated as part of a case-control study of moderate to severe acute diarrhea among children[24]. Among examined isolates prevalence of *aggR* and *aap* genes were 69.40% and 71.90% respectively and belonged to four phylogenetic groups A, B1, B2 and D. In the current study among 142 isolates nine *eaeA*+ isolates were detected which considered as atypical EPEC pathotype. Strains of EPEC are a well-known cause of diarrhoea particularly in infants and young children in less developed countries[26]. The results of an investigation on children with and without diarrhea in three Iranian Provinces, Tehran, Ilam and Mazandaran as a reservoir for intimin gene positive *E. coli* types showed that 40.5% and 20.0% of children with and without diarrhea harbored *eaeA* gene respectively[12]. On another study on 1610 *E. coli* isolates from patient age ranged from a few days to 98 years, 8.9% isolates were positive for EPEC and 4.8% positive for EAggEC pathotypes and 17 isolates were positive for both pathogens[27]. Phylogenetic analysis of DEC showed that EPEC strains were clustered mostly in groups B1, B2 and E[23]. The EIEC coding genetic marker (*ipaH*+) was a low frequency gene in the diarrheic isolates (4.92%). Similar reports showed that two isolates of the *E. coli* isolates from diarrheic children were positive for *ipaH* gene[22,28]. EIEC outbreaks are usually food or water borne; however, person-to-person transmission has also been reported[6]. This pathotype is extremely rare in southeast of Asia[8]. In a study, presence of the invasion-associated locus (*ial*) of the invasion plasmid was reported in 5% of children under two years old[29]. In Costa Rica, distribution of EIEC pathotype in each phylo-group indicated that isolates fell into A, B1 and D groups, whereas according to the results *ipaH* gene belonged to D phylogenetic group[3]. Phylogenetic analyses have shown that DEC strains fall into A, B1 and D phylo-groups[2]. None of the isolates possessed *stx1* and *stx2* genes and were not categorized as EHEC. This pathotype cannot be considered a main cause of childhood diarrhea in this region. These results are in accordance with the previous studies which were done on Thailand and Myanmar[30,31].

In conclusion ETEC and EAggEC were recovered at high

rates from children with diarrhea, indicating a wide spread of these pathotypes in the study population. It is maybe that the proportion of *E. coli* pathotypes difference according to the geographic region. The PCR assay can facilitate epidemiologic surveillance of DEC contamination. It is may also be used in epidemiologic surveillance of water for human consumption and food samples for *E. coli* contamination. Moreover, these contaminations can be transmitted from adults to children. Detection of epidemiological information may contribute to the prevention, including vaccines and control of infectious diarrhea in children.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

This work was supported by a grant (No: 111.IDT.91) from Research Center for Tropical and Infectious Diseases, Kerman University of Medical Sciences, Kerman, Iran. The authors are thankful to Dr. Eric Oswald (Ecole Nationale Vétérinaire Toulouse, France) for providing the reference strains.

References

- [1] Franciczek R, Sobieszczanska B, Turniak M, Kasprzykowska U, Krzyzanowska B, Jermakow K, et al. ESBL-producing *Escherichia coli* isolated from children with acute diarrhea—antimicrobial susceptibility, adherence patterns and phylogenetic background. *Adv Clin Exp Med* 2012; **21**: 187–192.
- [2] Mokracka J, Koczura R, Jablonska L, Kaznowski A. Phylogenetic groups, virulence genes and quinolone resistance of integron-bearing *Escherichia coli* strains isolated from a wastewater treatment plant. *Antonie Leeuwenhoek* 2011; **99**: 817–824.
- [3] Perez C, Gomez-Duarte OG, Arias ML. Diarrheagenic *Escherichia coli* in children from Costa Rica. *Am J Trop Med Hyg* 2010; **83**: 292–297.
- [4] Aranda KRS, Fagundes-Neto U, Scaletsky IC. Evaluation of multiplex PCRs for diagnosis of infection with diarrheagenic *Escherichia coli* and *Shigella* spp. *J Clin Microbiol* 2004; **42**: 5849–5853.
- [5] Mendez-Arancibia E, Vargas M, Soto S, Ruiz J, Kahigwa E, Schellenberg D, et al. Prevalence of different virulence factors and biofilm production in enteroaggregative *Escherichia coli* isolates causing diarrhea in children in Ifakara (Tanzania). *Am J Trop Med Hyg* 2008; **78**: 985–989.

- [6] Ahmed AM, Miyoshi S, Shinoda S, Shimamoto T. Molecular characterization of a multidrug-resistant strain of enteroinvasive *Escherichia coli* O164 isolated in Japan. *J Med Microbiol* 2005; **54**: 273–278.
- [7] Liebchen A, Benz I, Mellmann A, Karch H, Gomes TAT, Yamamoto D, et al. Characterization of *Escherichia coli* strains isolated from patients with diarrhea in Sao Paulo, Brazil: identification of intermediate virulence factor profiles by multiplex PCR. *J Clin Microbiol* 2011; **49**: 2274–2278.
- [8] Samosornsuk S, Chaicumpa W, von Seidlein L, Clemens JD, Sethabutr O. Using real time PCR to detect shigellosis: *ipaH* detection in Kaeng-Khoi District, Saraburi Province, Thailand. *Thammasat Int J Sci Technol* 2007; **12**: 52–57.
- [9] Sobieszczanska B, Kasprzykowska U, Turniak M, Maciejewski H, Franciczek R, Duda-Madej A. Virulence genes profiles and phylogenetic origin of *Escherichia coli* from acute and chronic intestinal diseases revealed by comparative genomic hybridization microarray. *Pol J Microbiol* 2012; **61**: 105–110.
- [10] Aslani MM, Bouzari S. Characterization of virulence genes of non-O157 Shiga toxin-producing *Escherichia coli* isolates from two provinces of Iran. *Jpn J Infect Dis* 2009; **62**: 16–19.
- [11] Konno T, Yatsuyanagi J, Takahashi S, Kumagai Y, Wada E, Chiba M, et al. Isolation and identification of *Escherichia albertii* from a patient in an outbreak of gastroenteritis. *Jpn J Infect Dis* 2012; **65**: 203–207.
- [12] Alikhani MY, Mirsalehian A, Aslani MM. Detection of typical and atypical enteropathogenic *Escherichia coli* (EPEC) in Iranian children with and without diarrhea. *J Med Microbiol* 2006; **55**: 1159–1163.
- [13] Nweze EI. Aetiology of diarrhoea and virulence properties of diarrhoeagenic *Escherichia coli* among patients and healthy subjects in southeast Nigeria. *J Health Popul Nutr* 2010; **28**: 245–252.
- [14] Carlos C, Pires MM, Stoppe NC, Hachich EM, Sato MIZ, Gomes TAT, et al. *Escherichia coli* phylogenetic group determination and its application in the identification of the major animal source of fecal contamination. *BMC Microbiol* 2010; doi: 10.1186/1471-2180-10-161.
- [15] China B, Pirson V, Mainil J. Typing of bovine attaching and effacing *Escherichia coli* by multiplex *in vitro* amplification of virulence-associated genes. *Appl Environ Microbiol* 1996; **62**: 3462–3465.
- [16] Cerna JF, Nataro JP, Estrada-Garcia T. Multiplex PCR for detection of three plasmid-borne genes of enteroaggregative *Escherichia coli* strains. *J Clin Microbiol* 2003; **41**: 2138–2140.
- [17] Clermont O, Bonacorsi S, Bingen E. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* 2000; **66**: 4555–4558.
- [18] Ghanbarpour R, Akhtardanesh B, Afsahi E, Sookhtanloo S. Molecular characterization of *Escherichia coli* pathotypes from diarrheic and healthy dogs. *Online J Vet Res* 2010; **14**: 316–324.
- [19] Jafari A, Aslani MM, Bouzari S. *Escherichia coli*: a brief review of diarrheagenic pathotypes and their role in diarrheal diseases in Iran. *Iran J Microbiol* 2012; **4**: 102–117.
- [20] Qadri F, Svennerholm AM, Faruque ASG, Sack RB. Enterotoxigenic *Escherichia coli* in developing countries: epidemiology, microbiology, clinical features, treatment, and prevention. *Clin Microbiol Rev* 2005; **18**: 465–483.
- [21] Jafari F, Hamidian M, Rezadehbashi M, Doyle M, Salmanzadeh-Ahrabi S, Derakhshan F, et al. Prevalence and antimicrobial resistance of diarrheagenic *Escherichia coli* and *Shigella* species associated with acute diarrhea in Tehran, Iran. *Can J Infect Dis Med Microbiol* 2009; **20**: e56–e62.
- [22] Vilchez S, Reyes D, Paniagua M, Bucardo F, Mollby R, Weintraub A. Prevalence of diarrhoeagenic *Escherichia coli* in children from Leon, Nicaragua. *J Med Microbiol* 2009; **58**: 630–637.
- [23] Escobar-Paramo P, Clermont O, Blanc-Potard AB, Bui H, Le Bouguenec C, Denamur E. A specific genetic background is required for acquisition and expression of virulence factors in *Escherichia coli*. *Mol Biol Evol* 2004; **21**: 1085–1094.
- [24] Boisen N, Scheutz F, Rasko DA, Redman JC, Persson S, Simon J, et al. Genomic characterization of enteroaggregative *Escherichia coli* from children in Mali. *J Infect Dis* 2012; **205**: 431–444.
- [25] Usein CR, Tatu-Chitoiu D, Ciontea S, Condei M, Damian M. *Escherichia coli* pathotypes associated with diarrhea in Romanian children younger than 5 years of age. *Jpn J Infect Dis* 2009; **62**: 289–293.
- [26] Clarke SC, Haigh RD, Freestone PPE, Williams PH. Virulence of enteropathogenic *Escherichia coli*, a global pathogen. *Clin Microbiol Rev* 2003; **16**: 365–378.
- [27] Hardegen C, Messler S, Henrich B, Pfeffer K, Wurthner J, MacKenzie CR. A set of novel multiplex Taqman real-time PCRs for the detection of diarrhoeagenic *Escherichia coli* and its use in determining the prevalence of EPEC and EAEC in a university hospital. *Ann Clin Microbiol Antimicrob* 2010; doi: 10.1186/1476-0711-9-5.
- [28] Hien BTT, Scheutz F, Cam PD, Serichantalergs O, Huong TT, Thu TM, et al. Diarrheagenic *Escherichia coli* and *Shigella* strains isolated from children in a hospital case-control study in Hanoi, Vietnam. *J Clin Microbiol* 2008; **46**: 996–1004.
- [29] Estrada-Garcia T, Lopez-Saucedo C, Thompson-Bonilla R, Abonce M, Lopez-Hernandez D, Santos JI, et al. Association of diarrheagenic *Escherichia coli* pathotypes with infection and diarrhea among Mexican children and association of atypical enteropathogenic *E. coli* with acute diarrhea. *J Clin Microbiol* 2009; **47**: 93–98.
- [30] Ratchtrachenchai OA, Subpasu S, Hayashi H, Ba-Thein W. Prevalence of childhood diarrhoea-associated *Escherichia coli* in Thailand. *J Med Microbiol* 2004; **53**: 237–243.
- [31] Takahashi E, Sultan Z, Shimada S, Aung WW, Nyein MM, Oo KN, et al. Studies on diarrheagenic *Escherichia coli* isolated from children with diarrhea in Myanmar. *Microbiol Immunol* 2008; **52**: 2–8.